



## Standard Practice for Algal Growth Potential Testing with *Pseudokirchneriella subcapitata*<sup>1,2</sup>

This standard is issued under the fixed designation D 3978; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ε) indicates an editorial change since the last revision or reapproval.

### INTRODUCTION

Algae are natural inhabitants of surface waters and are found in almost every water environment that is exposed to sunlight. The algae contribute to self purification (both organic and inorganic) of streams and lakes and are necessary as food for fish and fish food organisms. When large amounts of nutrients are available, excessive growths referred to as “blooms” can occur. Some algal blooms release substances toxic to fish, birds, domestic animals, and other alga. When nutrients are exhausted, the growth of algae and production of oxygen by photosynthesis decreases. The respiration of bacteria decomposing the large quantity of algal cells can deplete dissolved oxygen to the extent that fish and other oxygen consumers die. Both the abundance and composition of algae are related to water quality, with algal growth primarily influenced by the availability of nutrients.

The presence of indigenous algae in a water sample suggests that they are the most fit to survive in the environment from which the sample was taken. The indigenous algae should produce biomass until limited from further growth by some essential nutrient. If the indigenous algal production is limited from further growth by an essential nutrient, the laboratory test alga cultured in a noncompetitive environment and responding to the same limiting nutrient will produce parallel maximum yield growth responses. Generally, indigenous phytoplankton bioassays are not necessary unless there is strong evidence of the presence of long-term sublethal toxicants to which indigenous populations might have developed tolerance (1)<sup>3</sup>.

A single-indigenous algal species, dominant at the time of sampling, may not be more indicative of natural conditions than a laboratory species that is not indigenous to the system. The dynamics of natural phytoplankton blooms, in which the dominant algal species changes throughout the growth season, makes it quite certain that even if the indigenous algal isolate was dominant at the time of collection, many other species will dominate the standing crop as the season progresses.

When comparing algal growth potentials from a number of widely different water sources there are advantages in using a single species of alga. The alga to be used must be readily available and its growth measured easily and accurately. It must also respond to growth substances uniformly. Because some algae are capable of concentrating certain nutrients in excess of their present need when they are grown in media with surplus nutrients, this factor must be taken into account in selecting the culture media and in determining the type and amount of algae to use. (2) showed that a blue-green algae *Microcystis aeruginosa*, cultured in a low-nitrogen concentration medium, would not grow when transferred to medium lacking nitrogen. However, when the alga was cultured in medium containing four times as much nitrogen it was able to increase growth two-fold after transfer into nitrogen-free medium. A green alga *Pseudokirchneriella subcapitata* (formerly known as *Selenastrum capricornutum*, gave a similar response. In an analogous experiment with phosphorus, both organisms increased four-fold when transferred to medium lacking phosphorus. However, if algae are cultured in relatively dilute medium as recommended in the Algal Assay Procedure: Bottle Test (3) for culturing *Pseudokirchneriella subcapitata*, disclosed no significant further growth in medium lacking nitrogen or phosphorus when these were transferred from the initial medium over a wide range of inoculum sizes (4).

There are several methods available for determining algal growth. Measurements of optical density, oxygen production, carbon dioxide uptake, microscopical cell counts, and gravimetric cell mass determinations have been used, but often lack sensitivity when the number of cells is low.