



Standard Guide for Conducting Static and Flow-Through Acute Toxicity Tests With Mysids From the West Coast of the United States¹

This standard is issued under the fixed designation E 1463; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ε) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This guide describes procedures for obtaining data concerning the adverse effects of a test material (not food) added to marine and estuarine waters on certain species of marine and estuarine mysids during 96 h of continuous exposure. Juvenile mysids used in these tests are taken from cultures shortly after release from the brood and exposed to varying concentrations of a toxicant in static or flow-through conditions. These procedures will be useful for conducting toxicity tests with other species of mysids, although modifications might be necessary.

1.2 Modifications of these procedures might be justified by special needs or circumstances. Although using appropriate procedures is more important than following prescribed procedures, results of tests conducted using unusual procedures are not likely to be comparable to results of many other tests. Comparisons of results obtained using modified and unmodified versions of these procedures might provide useful information concerning new concepts and procedures for conducting acute tests with other species of mysids.

1.3 The procedures given in this guide are applicable to most chemicals, either individually or in formulations, commercial products, and known or unknown mixtures. With appropriate modifications these procedures can be used to conduct acute tests on factors such as temperature, salinity, and dissolved oxygen. These procedures can also be used to assess the toxicity of potentially toxic discharges such as municipal wastes, oil drilling fluids, produced water from oil well production, and other types of industrial wastes.

1.4 Results of acute toxicity tests with toxicants experimentally added to salt and estuarine waters should usually be reported in terms of a LC50 (median lethal concentration).

1.5 This guide is arranged as follows:

	Section
Referenced Documents	2
Terminology	3

Summary of Guide	4
Significance and Use	5
Apparatus	6
Facilities	6.1
Construction Materials	6.2
Metering Systems	6.3
Test Chambers	6.4
Cleaning	6.5
Acceptability	6.6
Safety Precautions	7
Dilution Water	8
Requirements	8.1
Source	8.2
Treatment	8.3
Characterization	8.4
Test Material	9
General	9.1
Stock Solution	9.2
Test Concentrations	9.3
Test Organisms	10
Species	10.1
Age	10.2
Source	10.3
Brood Stock	10.4
Food	10.5
Handling	10.6
Harvesting Young	10.7
Quality	10.8
Procedure	11
Experimental Design	11.1
Dissolved Oxygen	11.2
Temperature	11.3
Loading	11.4
Salinity	11.5
Light	11.6
Beginning of Test	11.7
Feeding	11.8
Duration of Test	11.9
Biological Data	11.10
Other Measurements	11.11
Analytical Methodology	12
Acceptability of Test	13
Interpretation of Results	14
Report	15
Appendixes	
<i>Holmesimysis costata</i>	X1
<i>Neomysis mercedis</i>	X2

1.6 The values stated in SI units are to be regarded as the standard.

¹ This guide is under the jurisdiction of ASTM Committee E47 on Biological Effects and Environmental Fate and is the direct responsibility of Subcommittee E47.01 on Aquatic Assessment and Toxicology.

Current edition approved April 1, 2004. Published April 2004. Originally approved in 1992. Last previous edition approved in 1998 as E 1463 – 92 (1998).